

REVIEW

An Obstructive Sleep Apnea - A Novel Public Health Threat

Sona GRESOVA¹, Martina GABOROVA¹, Judita STIMMELOVA¹, Igor PEREGRIM¹, Pavol SVORC¹, Viliam DONIC¹, Maria PALLAYOVA¹

¹Department of Human Physiology, Pavol Jozef Safarik University Faculty of Medicine, Kosice, Slovak Republic

Received January 16, 2023

Accepted April 18, 2023

Summary

In patients with obstructive sleep apnea (OSA) during obstructive events, episodes of hypoxia and hypercapnia may modulate the autonomic nervous system (ANS) by increasing sympathetic tone and irritability, which contributes to sympathovagal imbalance and ultimately dysautonomia. Because OSA can alter ANS function through biochemical changes, we can assume that heart rate variability (HRV) will be altered in patients with OSA. Most studies show that in both the time and frequency domains, patients with OSA have higher sympathetic components and lower parasympathetic dominance than healthy controls. These results confirm autonomic dysfunction in these patients, but also provide new therapeutic directions. Respiratory methods that modulate ANS, e.g., cardiorespiratory biofeedback, could be beneficial for these patients. Heart rate variability assessment can be used as a tool to evaluate the effectiveness of OSA treatment due to its association with autonomic impairment.

Key words

Autonomic nervous system • Heart rate variability • Obstructive Sleep apnea • Sleep

Corresponding author

Sona Gresova, Dpt. Physiology, Faculty of Medicine, Univ. PJ Safarik, Tr. SNP 1, 040001 Kosice, Slovak Republic. E-mail: sona.gresova@upjs.sk

Introduction

Obstructive sleep apnea (OSA) is defined as the absence of ventilation lasting for at least 10 seconds, during which the activity of the respiratory center is

maintained so the respiratory effort remains. This phenomenon is caused by the paradoxical movement of the chest and diaphragm that occurs during an obstructive sleep apnea episode. This is a recurrent collapse of the upper airways during sleep, when transient obstruction during snoring is caused by weakening and "sucking" of the oropharyngeal walls with the interruption or restriction of air permeability and a consequent decrease in blood oxygen saturation. Thus, transient hypoxemia occurs with sleep disturbance and fragmentation, resulting in excessive daytime sleepiness. Due to vagal activity, the heart rate and blood pressure decrease during the apnea pause. A decrease in pO₂ and increase in pCO₂ activate chemoreceptors in the carotid sinus, aortic arch and spinal cord. This activation leads to a reflex increase in ventilation, causing microarousals and increased muscle tone of the upper respiratory tract, which subsequently opens. As a result of sympathetic activation, persons with OSA hyperventilate, have tachycardia and their blood pressure increases. Increased sympathetic activity leads to vasoconstriction and increased peripheral vascular resistance in the systemic and pulmonary systems. After falling asleep again, this whole cycle is repeated. These periods are accompanied by snoring of varying intensity (snoring can also be a predictor of OSA).

Heart rate variability (HRV) can be used to evaluate the integrity of the autonomic nervous system (ANS) in these patients. A lower HRV is observed in several cardiovascular and respiratory diseases, such as hypertension and chronic obstructive pulmonary disease (2), indicating the abnormal adaptability of ANS in these

patients. As OSA can alter the ANS function through biochemical changes, we can assume that HRV will be altered in patients with obstructive sleep apnea (3).

Previous studies have yielded promising results showing that heart rate variability analysis can be a predictor of sleep apnea, reaching a sensitivity and specificity of approximately 70–90 % [1]. However, it should be emphasized that these studies were limited to small groups of patients and were primarily based on qualitative rather than quantitative analyses. The primary goal of the present review was to describe changes in the autonomic nervous system in patients with OSA.

Obstructive sleep apnea

Obstructive sleep apnea is the most common type of sleep apnea, accounting for up to 80 % of all cases of sleep apnea syndrome. According to a 2016 study [2], the overall prevalence of OSA (defined by an apnea-hypopnea index/AHI ≥ 5 respiratory events/h) ranged from 9–38 % in the general population and was higher in men. The OSA prevalence increased with age, reaching up to 90 % in men aged 60–85 years and 78 % in women of a similar age [2]. When defined by the AHI ≥ 15 respiratory events/h, the prevalence of OSA ranged from 6–17 % in the adult population (49 % were elderly) [2]. The awareness of doctors about OSA and its complications plays a key role in providing necessary healthcare. In places where these disorders are given increased attention, patients are sent for examinations for suspected OSA much more often [2]. This leads to a locally increased percentage of diagnosed patients [2]. Obstructive sleep apnea occurs in all age categories, but the most common period of symptoms is in the 4th and 5th decade of life. During this period, the symptoms are also the most significant [2].

The incidence of OSA is higher in obese patients, in type 2 diabetes mellitus and in other cardiovascular disorders which include coronary artery disease, heart failure, cardiac arrhythmias, stroke, atherosclerosis and myocardial infarction. Research by the Seventh Joint National Committee shows that OSA is the cause of secondary hypertension in 37–56 % of cases and the prevalence rate of OSA in patients with resistant hypertension is estimated at 70–83 % [3]. In the pediatric population, the incidence of OSA ranges from 1.2–5.7 % and the risk of developing OSA in obese children is estimated at 36 % [4]. It is reported that in more than 85 % of patients with OSA, the symptoms have never

been clinically diagnosed [5]. The incidence of snoring itself is much higher, affecting one in three adults, but it is a habitual snoring that is not accompanied by clinical signs of the disease. We also refer to this type of snoring as simple benign snoring [6].

Pathogenesis of obstructive sleep apnea

Obstructive sleep apnea is characterized by recurrent episodes of upper airway collapse and obstruction during sleep due to defects in the pharyngeal structure and neuromuscular control [7]. Important components include abnormalities in the upper airway anatomy, the ability of the upper airway dilator muscles to respond to the challenge of breathing during sleep, the tendency to wake up after increased respiratory effort, the stability of the respiratory center and the potential for lung volume changes to influence these factors. Also, OSA is often associated with changes in upper airway anatomy; structural changes, including tonsil hypertrophy, retrognathion and variations in craniofacial structures are all associated with an increased risk of sleep apnea as they contribute to increased upper airway collapse. Ethnic differences in craniofacial traits are one of the potential explanations for the observed differences in the prevalence and severity of OSA for a given level of obesity. Studies in awake patients using CT and MRI have shown increased adipose tissue deposition and submucosal oedema in the lateral walls of the pharynx, both of which contribute to narrowing of the pharyngeal lumen and may predispose these individuals to obstruction during sleep [8]. Based on the presence of anatomical changes in the upper airways in patients with OSA, structural or mechanical changes are considered to be the primary determinant of upper airway obstruction during sleep [8]. Studies have shown that structural changes in the lateral walls of the pharynx and tongue arise on a family basis, suggesting a role for genetics in the development of OSA. In addition, experimental data obtained in the absence of neuromuscular activity show narrowing of the pharyngeal space and increased Pcrit (passive pharyngeal critical closing pressure) values in individuals with OSA. In addition, obesity, jaw position, acromegaly, tonsillar hypertrophy and the smaller bone envelope surrounding the pharynx have been shown to increase pharyngeal collapse [8]. Upper airway structural differences differentiate OSA patients from healthy subjects. In addition, when neuromuscular mechanisms weaken during sleep, they may predispose them to upper

airway obstruction [8].

Obesity is another serious risk factor for OSA. It is associated with an increased neck circumference and increased amounts of peripheral fat, which can narrow and compress the upper airways. Increased parapharyngeal fat also correlates with the increasing severity of sleep apnea. Studies examining resected uvular tissue have revealed greater amounts of submucosal adipose tissue in patients with OSA. The compressive effects of adipose tissue deposited around the pharynx may therefore increase the collapse of the upper airways and possibly antagonize the effects of the expansion muscles that maintain their patency. Obesity also increases pharyngeal collapse through a reduction in lung volumes, especially by reducing functional residual capacity (FRC), which deepens with the onset of sleep. Decreased FRC increases pharyngeal collapse by reducing tracheal traction on the pharyngeal segment. Conversely, an increase in lung volumes leads to an increase in tracheal traction and stabilization of the upper airways during inspiration. In patients with OSA, an increase in lung volumes has been shown to reduce continuous positive airway pressure (CPAP) requirements and the severity of OSA [9].

Neuromuscular factors

Anatomical mechanical loads on the upper airways may not always lead to pharyngeal collapse during sleep. Women have a smaller pharynx and oropharyngeal junction than men, but still have a lower prevalence of OSA. In addition, measurements of Perit under low neuromuscular activity conditions, which reflect the mechanical loads on the upper airways, show significant overlap between OSA patients and controls. Thus, non-structural (i.e., neuromuscular) factors must also play a role in upper airway protection. Changes in upper airway neuromuscular activity during sleep were originally described by Remmers *et al.*, who showed that genioglossal electromyogram (EMG_{gg}) activity was reduced at the onset of apnea and increased at arousal when airway patency was restored. Subsequently, upper airway obstruction has been found to trigger various neuromuscular responses that restore upper airway patency by activating muscles that dilate and lengthen the airways. During the respiratory cycle, various pharyngeal muscle groups are important, which ensure stabilization (tonic activity, e.g., m. tensor palatini) and airway dilatation during inspiration (phase activity, e.g., m. genioglossus). The resulting pharyngeal motility is

modulated by a number of factors, which include waking and sleep-dependent mechanisms, local mechanoreceptor responses to vacuum and respiratory control mechanisms [10]. During wakefulness, an increased activity of m. genioglossus and m. tensor palatini was observed in patients with OSA, which can be significantly reduced by applying positive pressure. On the contrary, the control group of healthy individuals showed lower activity of these muscles, which did not decrease further after the application of CPAP. These findings suggest that the increased upper airway dilatation muscle activity compensates for anatomically narrower upper airways in patients with OSA. Decreased upper airway muscle activity with the onset of serotonergic, cholinergic, noradrenergic and histaminergic pathways may lead to their obstruction [11].

Pressure-sensing mechanisms play an important role in modulating upper airway neuromuscular activity during wakefulness and sleep. The negative pressure in the upper respiratory tract serves to stabilize them during inhalation. The vacuum reflex is primarily mediated by mechanoreceptors in the pharynx. This argument is supported by the fact that there is a relationship between EMG_{gg} (m. genioglossus electromyography) and pharyngeal pressure independent of the central respiration generated in the brainstem. Local anesthesia on the pharyngeal mucosa weakens this relationship, leading to increased obstructive apnea and hypopnea, loud snoring and/or prolonged apnea episodes in healthy subjects, implying that the pharyngeal vacuum stabilizes upper airway patency during inhalation [11].

Microarousals

Waking from sleep at the end of hypopnea or apnea has long been considered an important protective mechanism for reopening the airways [12]. However, work by Younes [12] provided insights into the functional role of waking from sleep in OSA and questioned the notion that it was a necessary mechanism for reopening the airways. When examining the response to experimentally-induced transient airway depression (CPAP) in patients with OSA, Younes noted that inspiratory flow increased in 22 % of cases before awakening and resumed in 17 % of cases without awakening [12]. Jordan *et al.* conducted a study on the mechanisms underlying the resumption of airflow without awakening patients [13]. A transient decrease in pressure for up to 5 minutes resulted in an increase in genioglossus muscle activity and changes in the work

cycle. These compensatory responses were similar between OSA patients and healthy individuals. However, compared to healthy patients receiving stimuli of a similar magnitude, patients with OSA without increased cortical activity had a lower ability to restore ventilation [10]. Findings that patients with OSA are able to restore ventilation at respiratory loads without cortical arousal for at least some time increase the likelihood that some patients are able to maintain clear airways during sleep if they remain in sleep long enough to restore compensatory mechanisms [10]. Combinations of stimuli, such as carbon dioxide and vacuum, can activate the dilatation muscles of the upper respiratory tract [10]. Delaying the awakening, which would allow for the sufficient accumulation of respiratory stimuli to restore pharyngeal patency, could thus be beneficial for patients. Procedures to prevent awakenings from sleep (increasing the individual awakening threshold) are likely to be the most effective in patients who wake up easily during sleep after sleep exercise. However, raising the awakening threshold in patients who already have a high value may be detrimental due to the possible worsening of blood gas abnormalities [10]. Most of the available evidence suggests that the level of pleural pressure generated by respiratory effort regardless of stimulus (e.g., hypoxia, hypercapnia and respiratory load) is likely to be a key trigger to induce non-REM sleep awakening [10]. Experimentally, the awakening threshold is measured as the minimum oesophageal pressure (or pressure at the level of the epiglottis that is thought to be similar during airway occlusion) generated by the breath before awakening during respiratory load or occlusion. Although there is considerable interindividual variability, patients with OSA tend to have a worsened response to micro-arousal (greater negative pressure or higher awakening threshold is required) than controls. CPAP treatment tends to lower the awakening threshold [10]. These findings suggest that OSA itself (e.g., sleep fragmentation, hypoxia and recurrent airway obstruction) is responsible for impaired awakening responses in OSA patients and not an inherent abnormality of the awakening threshold. Awakening from sleep is associated with increased upper airway dilator activity at an equivalent level of negative pharyngeal pressure during sleep and with a rapid ventilatory response. Although these changes are beneficial in rapidly restoring airflow and reversing oxygen desaturation/hypercapnia, they can also destabilize respiration and contribute to the severity of apnea [10].

Besides epigenetic factors, genetic predisposition may play an important role in the pathogenesis of OSA. Studies have clearly shown a familial occurrence in the development of obstructive sleep apnea [14]. This finding applies to both obese and non-obese patients with OSA. Studies using linkage analysis have provided an initial insight into the possible linkages between specific regions of the genome and the pathogenesis of OSA. Anatomy (obesity, craniofacial structure) clearly has a genetic basis [14]. In addition, features such as upper airway soft tissue structure size, respiratory control abnormalities and respiratory responses to sleep resistance may also have a genetic background [14].

Heart rate variability in obstructive sleep apnea

Recurrent episodes of obstructive sleep apnea lead to periodic hypoxemia, hypercapnia and arousal from sleep, which stimulates peripheral and central chemoreceptors. They cause cyclic oscillations in the activity of the sympathetic and parasympathetic nervous systems, which cause oscillations in blood pressure and heart rate with each apnea-hyperpnoea cycle. Autonomic regulation is related to baroreceptors and chemoreceptors, which can perceive biochemical modifications in pCO₂ and pH. Hypoxia and hypercapnia during recurrent obstructive episodes are therefore considered to be a major cause of altered ANS. During obstructive events, these episodes of hypoxia and hypercapnia may modulate the ANS of patients with OSA by increasing sympathetic tone and irritability, which may contribute to sympathovagal imbalance and ultimately dysautonomia [15].

Several physiological factors derived from recurrent upper airway obstruction can lead to autonomic dysfunction and the cardiovascular comorbidities of OSA. Obstructive events increase negative intrathoracic pressure and hypoxia, which has adverse hemodynamic consequences, such as excessive sympathetic activity, enhanced reactive oxygen species (ROS) and vasoactive compensatory mechanisms [16]. Sympathetic activity and blood pressure are also determined by the duration of the obstructive episode and the severity of oxygen desaturation [17].

HRV assessment can be used as a tool to evaluate the effectiveness of treatment of patients with OSA due to its association with autonomic damage. In

frequency domain analysis, VLF and ULF are mainly influenced by SF circadian differences and therefore cannot be estimated by short-term analysis, only by a 24-hour Holter [18]. Elevated VLF levels may be associated with the increased sympathetic activity observed in OSA patients in some studies. Both components account for 95 % of total performance and the literature suggests that they have the highest predictive value because they are more strongly associated with overall health than the high-frequency components. There are several limitations in studies looking at HRV changes in patients with OSA. In HRV research, the effect of the menstrual cycle on autonomic cardiac modulation is less studied in women. Although OSA is more common in men than in women, further investigation of sex differences in HRV in these patients could clarify whether hormones are also involved in this sleep-related breathing disorder [19]. Several studies also include participants with associated comorbidities and patients taking medications that may affect HRV outcomes. Additional limitations include different data length, device and record status. These differences between the studies affect the accuracy of HRV analysis and especially the comparison of their results [20].

The values measured in the low-frequency LF band are mainly derived from sympathetic activity, but they also reflect the activity of the parasympathetic nervous system to a lesser extent. The vast majority of studies have shown that healthy control groups have significantly lower LF values than patients with OSA. However, the results in the literature vary. There are several studies that did not confirm statistically significant differences in the LF parameter and one study in which the control group had significantly higher LF values than the observed group of patients with OSA [17].

The HF high frequency band is a component of HRV that represents vagal cardiac activity. High HF levels are considered a predictor of good health. Studies have shown that patients with OSA have significantly lower HF levels than healthy controls, indicating a lower parasympathetic predominance and an increased risk of cardiovascular problems. Increased HF performance in patients with severe OSA may be associated with an increase in vagal tone, presumably due to more frequent and intensive Mueller maneuvers. A large change in the respiratory model of sick patients may affect HF performance, as the high frequency band is also known to be affected by respiration itself [17].

The low frequency to high frequency ratio HRV (LF/HF) represents the sympathovagal balance index. High LF/HF values indicate a predominance of sympathetic activity and dysautonomia. Most studies have shown elevated LF/HF levels in OSA patients compared to healthy controls. A higher proportion of sympathetic tone is probably related to recurrent apnea events. Relative hypoxemia acts on central chemoreceptors and thus increases sympathetic activity. Repeated arousal from sleep can also cause relatively high sympathetic activity and decreased vagal tone. A retrospective study of patients with varying severities of obstructive sleep apnea found that the arousal index is a stronger contributing factor to cardiac autonomic dysfunction in patients with OSA than the apnea-hypopnea index (AHI) [21]. Elevated values of the LF/HF standard deviation indicate that the changes in sympathetic and parasympathetic activity during the night are not identical. This may be due to less common apnea events.

Changes in the stage of sleep are also an important factor. REM sleep, which usually occurs at 90-minute intervals, is associated with a significantly higher number of apnea episodes in patients with OSA. The long 24-hour data collection is considered the "gold standard" because it involves changes in the body's metabolism and circadian rhythm. All of these selected studies also used AHI as a classification of the severity of sleep apnea, but this index does not fully correlate with cardiovascular risk in adults. The coherent study [22] showed that the prediction of cardiovascular outcomes can be independent of AHI. AHI represents only the number of apnea and hypopnea events per hour of sleep [22], which documents the frequency of these events, but not, for example, the duration or magnitude of each event. The study by Hietakoste *et al.*, in 2020 [23], provides valuable insight into the cardiovascular stress associated with apnea and hypopnea. The results show that higher ultrashort HRV and greater variation between RR intervals within and after a respiratory episode are more strongly associated with longer respiratory event duration, apnea and male gender. Although some of these selected studies compare data between AHI and HRV, more studies are needed to clarify whether the current definition of hypopnea and sleep apnea severity classification still apply as a predictor of cardiovascular risk in patients with OSA [17]. On the other hand, HRV can be used as a predictor of morbidity and mortality, especially SDNN, which is considered the "gold

standard" for medical stratification in 24-hour recordings [24].

Heart rate variability as an obstructive sleep apnea novel diagnostic tool

At the beginning of the investigation of obstructive sleep apnea, it was found that the events of apnea and hypopnea are accompanied by concomitant cyclical changes in heart rate. The bradycardia/tachycardia pattern is closely related to the time course of apnea events. As a result, this model has been successfully used to detect sleep apnea in patients with clinical symptoms. Over the last 15 years, special devices have been developed to detect sleep apnea in the early stages of the disease by recording heart rate, snoring, oxygen saturation and body position. These devices have been successfully used to stratify the individual risk of patients suffering from sleep-disordered breathing before being referred for cardiorespiratory polysomnography. The sensitivity and specificity of these simplified four-channel recording systems have been evaluated in many studies providing results for specific systems. A recent systematic review summarized the results of these studies in a comparative review [25]. This review revealed the value and limitations of the systems with respect to the accepted criteria for the diagnosis of sleep apnea. It should be noted that satisfactory standardization of terms and diagnostic criteria for a transmissible diagnosis of sleep apnea has not been achieved.

In parallel with efforts to define evidence-based criteria for the simplified diagnosis of sleep apnea, the physiology causing cyclical changes in heart rate has been investigated. Studies of sympathetic nerve activity during sleep apnea have shown that sympathetic activation increases during apnea [26]. It is not entirely clear to what extent nerve activity reflects the sympathetic activation of the heart during apnea with effects on the heart rate itself. Sympathetic nerve activation in patients with sleep apnea even persists during the day and may contribute to an increased cardiovascular risk in these patients [27]. The study was validated in 2020, in which HRV served as a supplement to well-known cardiovascular risk factors such as aging, adiposity and sleep disorders during sleep, with an overall accuracy of 75.3 %. This suggests that adding heart rate variability to the severity classification may predict a real impact on physiological parameters that later lead to pathological changes [28].

Several studies have examined heart rate and HRV before [29] and during sleep apnea in order to improve the means for a simplified diagnosis of sleep-disordered breathing. At the same time, the work suggests that changes in HRV are caused by sleep apnea [30]. One of these studies uses sympathetic activity derived from the spectral components of HRV. A standard technique is introduced to derive sympathetic and parasympathetic tone indicators from HRV by calculating the spectral power in defined frequency bands. This technique has been shown to have some prediction of hypertension and myocardial infarction and is therefore currently used in many cardiovascular disorders. The low frequency component of HRV is usually associated with sympathetic activity. This study examined the spectral components in short 2-minute time segments in patients with OSA and compares them to undisturbed sleep in healthy controls. The study concluded that there was an increased sympathetic tone in respiratory disorders [30]. However, sleep-associated respiration represents a complex interaction between regular changes in the autonomic nervous system during sleep and specific changes associated with apnea. From a physical point of view, spectral analysis requires stationary signals. This means that the mean and standard deviation and moments of the higher order of the analyzed signal must remain the same for the entire investigation period. Sleep apnea with a highly dynamic HRV pattern is by no means a stationary process; therefore, the use of spectral analysis is limited. Due to these limitations, other authors are trying to use new techniques developed specifically for non-stationary data analysis. Time-frequency wave analysis is a sophisticated technique superior to spectral analysis because it is not restricted to limiting stationary requirements [1]. The problem with wave analysis is that the results are less intuitive, so another method needs to be used to select the best wave to detect sleep apnea. In a research study by Roche *et al.* from 2003, (1) they used the wave method on a large sample of patients and developed an optimal algorithm in order to identify the most suitable waves. The presented results prove that this method is effective in recognizing cyclic HRV specific for sleep apnea, because the pattern is not strictly periodic. The application of this method can improve the recognition of sleep apnea based on non-invasive simple electrocardiography (ECG) and heart rate recording [31]. Both studies confirm the finding that new methods can detect sleep apnea only from HRV changes to a good extent [31]. Thus, it seems that simplified recording

techniques can be used in combination with sophisticated analysis techniques to detect sleep disorders. The 2021 review study provides, among other things, an overview of the physiological basis for measuring heart rate and heart rate variability in wearable devices and summarizes some of the strengths and limitations of consumer devices [32]. The latest pacemakers already offer an optional indication of sleep apnea based on an analysis of past heart rate records stored in their memory. As a result, more cardiologists using long-term ECG recordings and using pacemaker heart rate extracts can identify patients with suspected sleep apnea [33]. The physiology behind the HRV changes is remarkable and several aspects of heart-lung interactions have been elucidated to date. Therefore, it is interesting to investigate the time course of sympathetic and parasympathetic activation during apnea and subsequent breaths [26]. The time course of activation of the autonomic nervous system has been described by sympathetic nerve activity, but the time course of parasympathetic activity has not been determined. It is quite likely that both sympathetic and parasympathetic activity increase at the end of each period of apnea and then decrease over a short period of compensatory hyperventilation. However, volumetric hyperventilation in healthy subjects was associated with a predominance of sympathetic disease, but changes in hemodynamics and sympathovagal balance were smaller in patients with heart failure and arterial hypertension [34].

In a 2019 study, Festic *et al.* [35] observed changes in HRV and nocturnal pulse oximetry in subjects with an oxygen desaturation index of less than 5 events/h and found that the episodic increase in HRV occurring with minimal oxygen desaturation in normal nocturnal oximetry is a strong independent predictor of OSA, despite the increased use of drugs to control the heart rate in patients with OSA. It is possible that some quantifiable methods such as nocturnal pulse oximetry and pulse monitoring may provide better predictive power given the severity of OSA [35].

Impact of obstructive sleep apnea treatment on heart rate variability

CPAP is still the gold standard treatment for

moderate to severe obstructive sleep apnea. CPAP treatment affected changes in heart rate variability in patients with OSA; these changes are more significant using frequency analysis compared to time analysis and they are also noticeable in medium and long periods of treatment with significant modifications in terms of decreased sympathovagal imbalance and increased vagal component (HF) [36]. In a meta-analysis by Guo *et al.* [37], the researchers point out that CPAP treatment for at least 1 month in patients with obstructive sleep apnea is likely to improve sympathovagal imbalance, which may be one mechanism by which CPAP could reduce the risk of cardiovascular disease in patients with OSA.

Kufof *et al.* [38] emphasize that heart rate variability parameters acutely improve regardless of gender and weight on the first night of CPAP use in patients with severe OSA and therefore CPAP treatment should not be delayed; conversely, patients with severe OSA should be informed that even a single night without CPAP results in changes in HRV that are adjusted by CPAP treatment.

Conclusion

In conclusion, OSA is a highly prevalent but also largely underdiagnosed condition. Early diagnosis and satisfactory treatment are critical to prevent numerous OSA sequelae. In order to improve the healthcare process in the public health sector, the liaison between primary care physicians and sleep specialists is needed. Considering the continually increasing prevalence of obesity and diabetes worldwide, we can expect the epidemic of OSA to become exacerbated in the near future.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

This publication is the result of the project implementation: “Design and implementation of advanced methods of lungs ventilation treatment and the diagnosis of viral pneumonia, including COVID-19 with the possibility of their rapid adoption”, ITMS2014+: 313011ASX1 supported by the Operational Programme Integrated Infrastructure, funded by the ERDF.

References

1. Roche F, Pichot V, Sforza E, Court-Fortune I, Duverney D, Costes F, et al. Predicting sleep apnoea syndrome from heart period: a time-frequency wavelet analysis. *Eur Respir J.* 2003;22:937-42. <https://doi.org/10.1183/09031936.03.00104902>

2. Senaratna CV, Perret JL, Lodge CJ, Lowe AJ, Campbell BE, Matheson MC, et al. Prevalence of obstructive sleep apnea in the general population: A systematic review. *Sleep Med Rev.* 2017;34:70-81. <https://doi.org/10.1016/j.smr.2016.07.002>
3. Pedrosa RP, Drager LF, Gonzaga CC, Sousa MG, de Paula LK, Amaro AC, et al. Obstructive sleep apnea: the most common secondary cause of hypertension associated with resistant hypertension. *Hypertension (Dallas, Tex:1979).* 2011;58(5):811-817. <https://doi.org/10.1161/HYPERTENSIONAHA.111.179788>
4. Brockbank JC. Update on pathophysiology and treatment of childhood obstructive sleep apnea syndrome. *Paediatr Respir Rev.* 2017;24:21-23. <https://doi.org/10.1016/j.prrv.2017.06.003>
5. Motamedi KK, McClary AC, Amedee RG. Obstructive sleep apnea: a growing problem. *Ochsner J.* 2009;9:149-153.
6. De Meyer MMD, Jacquet W, Vanderveken OM, Marks LAM. Systematic review of the different aspects of primary snoring. *Sleep Med Rev.* 2019;45:88-94. <https://doi.org/10.1016/j.smr.2019.03.001>
7. McGinley BM, Schwartz AR, Schneider H, Kirkness JP, Smith PL, Patil SP. Upper airway neuromuscular compensation during sleep is defective in obstructive sleep apnea. *J Appl Physiol (1985).* 2008;105(1):197-205. <https://doi.org/10.1152/jappphysiol.01214.2007>
8. Watanabe T, Isono S, Tanaka A, Tanzawa H, Nishino T. Contribution of body habitus and craniofacial characteristics to segmental closing pressures of the passive pharynx in patients with sleep-disordered breathing. *Am J Respir Crit Care Med.* 2002;165(2):260-265. <https://doi.org/10.1164/ajrccm.165.2.2009032>
9. Heinzer RC, Stanchina ML, Malhotra A, Jordan AS, Patel SR, Lo YL, et al. Effect of increased lung volume on sleep disordered breathing in patients with sleep apnoea. *Thorax.* 2006;61:435-439. <https://doi.org/10.1136/thx.2005.052084>
10. Eckert DJ, Malhotra A. Pathophysiology of adult obstructive sleep apnea. *Proc Am Thorac Soc.* 2008;5:144-153. <https://doi.org/10.1513/pats.200707-114MG>
11. Chamberlin NL, Eikermann M, Fassbender P, White DP, Malhotra A. Genioglossus premotoneurons and the negative pressure reflex in rats. *J Physiol.* 2007;579(Pt 2):515-526. <https://doi.org/10.1113/jphysiol.2006.121889>
12. Younes M. Role of arousals in the pathogenesis of obstructive sleep apnea. *Am J Respir Crit Care Med.* 2004;169:623-633. <https://doi.org/10.1164/rccm.200307-1023OC>
13. Jordan AS, Wellman A, Heinzer RC, Lo YL, Schory K, Dover L, et al. Mechanisms used to restore ventilation after partial upper airway collapse during sleep in humans. *Thorax.* 2007;62:861-7. <https://doi.org/10.1136/thx.2006.070300>
14. Larkin EK, Patel SR, Redline S, Mignot E, Elston RC, Hallmayer J. Apolipoprotein E and obstructive sleep apnea: evaluating whether a candidate gene explains a linkage peak. *Genet Epidemiol.* 2006;30:101-110. <https://doi.org/10.1002/gepi.20127>
15. Kheirandish-Gozal L, Bhattacharjee R, Gozal D. Autonomic alterations and endothelial dysfunction in pediatric obstructive sleep apnea. *Sleep Med.* 2010;11(7):714-20. <https://doi.org/10.1016/j.sleep.2009.12.013>
16. Khayat R, Pleister A. Consequences of Obstructive Sleep Apnea: Cardiovascular risk of obstructive sleep apnea and whether continuous positive airway pressure reduces that risk. *Sleep Med Clin.* 2016;11(3):273-286. <https://doi.org/10.1016/j.jsmc.2016.05.002>
17. Sequeira VCC, Bandeira PM, Azevedo JCM. Heart rate variability in adults with obstructive sleep apnea: a systematic review. *Sleep Sci.* 2019;12:214-221. <https://doi.org/10.5935/1984-0063.20190082>
18. Bauer A, Camm AJ, Cerutti S, Guzik P, Huikuri H, Lombardi F, et al. Reference values of heart rate variability. *Heart rhythm.* 2017;14:302-303. <https://doi.org/10.1016/j.hrthm.2016.12.015>
19. Tada Y, Yoshizaki T, Tomata Y, Yokoyama Y, Sunami A, Hida A, et al. The impact of menstrual cycle phases on cardiac autonomic nervous system activity: an observational study considering lifestyle (diet, physical activity, and sleep) among female college students. *J Nutr Sci Vitaminol (Tokyo).* 2017;63:249-255. <https://doi.org/10.3177/jnsv.63.249>
20. Taralov ZZ, Terziyski KV, Kostianev SS. Heart rate variability as a method for assessment of the autonomic nervous system and the adaptations to different physiological and pathological conditions. *Folia Med (Plovdiv).* 2015;57:173-180. <https://doi.org/10.1515/folmed-2015-0036>

21. Kim YS, Kim SY, Park DY, Wu HW, Hwang GS, Kim HJ. Clinical implication of heart rate variability in obstructive sleep apnea syndrome patients. *J Craniofac Surg.* 2015;26:1592-1595. <https://doi.org/10.1097/SCS.0000000000001782>
22. Campos-Rodriguez F, Martínez-García MA, Reyes-Nuñez N, Selma-Ferrer MJ, Punjabi NM, Farre R. Impact of different hypopnea definitions on obstructive sleep apnea severity and cardiovascular mortality risk in women and elderly individuals. *Sleep Med.* 2016;27-28:54-58. <https://doi.org/10.1016/j.sleep.2016.05.020>
23. Hietakoste S, Korkalainen H, Kainulainen S, Sillanmäki S, Nikkonen S, Myllymaa S, et al. Longer apneas and hypopneas are associated with greater ultra-short-term HRV in obstructive sleep apnea. *Sci Rep.* 2020;10:21556. <https://doi.org/10.1038/s41598-020-77780-x>
24. Shaffer F, Ginsberg JP. An overview of heart rate variability metrics and norms. *Frontiers in public health.* 2017;5:258. <https://doi.org/10.3389/fpubh.2017.00258>
25. Ucak S, Dissanayake HU, Sutherland K, de Chazal P, Cistulli PA. Heart rate variability and obstructive sleep apnea: Current perspectives and novel technologies. *J Sleep Res.* 2021;30(4):e13274. <https://doi.org/10.1111/jsr.13274>
26. Narkiewicz K, Somers VK. Sympathetic nerve activity in obstructive sleep apnoea. *Acta Physiol Scand.* 2003;177:385-390. <https://doi.org/10.1046/j.1365-201X.2003.01091.x>
27. Narkiewicz K, Pesek CA, Kato M, Phillips BG, Davison DE, Somers VK. Baroreflex control of sympathetic nerve activity and heart rate in obstructive sleep apnea. *Hypertension.* 1998;32:1039-1043. <https://doi.org/10.1161/01.HYP.32.6.1039>
28. Zhang L, Wu H, Zhang X, Wei X, Hou F, Ma Y. Sleep heart rate variability assists the automatic prediction of long-term cardiovascular outcomes. *Sleep Med.* 2020;67:217-224. <https://doi.org/10.1016/j.sleep.2019.11.1259>
29. Karimi Moridani M. A novel clinical method for detecting obstructive sleep apnea using of nonlinear mapping. *J Biomed Phys Eng.* 2022;12:31-34. <https://doi.org/10.31661/jbpe.v0i0.1211>
30. Dingli K, Assimakopoulos T, Wraith PK, Fietze I, Witt C, Douglas NJ. Spectral oscillations of RR intervals in sleep apnoea/hypopnoea syndrome patients. *Eur Respir J.* 2003;22:943-950. <https://doi.org/10.1183/09031936.03.00098002>
31. Penzel T. Is heart rate variability the simple solution to diagnose sleep apnoea? *Eur Respir J.* 2003;22(6):870-871. <https://doi.org/10.1183/09031936.03.00102003>
32. Lujan MR, Perez-Pozuelo I, Grandner MA. Past, Present, and future of multisensory wearable technology to monitor sleep and circadian rhythms. *Front Digit Health.* 2021;3:721919. <https://doi.org/10.3389/fdgh.2021.721919>
33. Penzel T, McNames J, Murray A, de Chazal P, Moody G, Raymond B. Systematic comparison of different algorithms for apnoea detection based on electrocardiogram recordings. *Med Biol Eng Comput.* 2002;40:402-407. <https://doi.org/10.1007/BF02345072>
34. Spiesshoefer J, Becker S, Tuleta I, Mohr M, Diller GP, Emdin M, et al. Impact of simulated hyperventilation and periodic breathing on sympatho-vagal balance and hemodynamics in patients with and without heart failure. *respiration.* 2019;98:482-494. <https://doi.org/10.1159/000502155>
35. Festic N, Zuberi M, Bansal V, Fredrickson P, Festic E. Correlation between oxygen saturation and pulse tracing patterns on overnight oximetry with normal desaturation index is an independent predictor of obstructive sleep apnea. *J Clin Sleep Med.* 2019;15(2):195-200. <https://doi.org/10.5664/jcsm.7614>
36. Correia FJ, Martins LEB, Barreto DM, Pithon KR. Repercussion of medium and long treatment period with continuous positive airways pressure therapy in heart rate variability of obstructive sleep apnea. *Sleep Sci.* 2019;12:110-115. <https://doi.org/10.5935/1984-0063.20190068>
37. Guo W, Lv T, She F, Miao G, Liu Y, He R, et al. The impact of continuous positive airway pressure on heart rate variability in obstructive sleep apnea patients during sleep: A meta-analysis. *Heart Lung.* 2018;47:516-524. <https://doi.org/10.1016/j.hrtlng.2018.05.019>
38. Kufoy E, Palma JA, Lopez J, Alegre M, Urrestarazu E, Artieda J, et al. Changes in the heart rate variability in patients with obstructive sleep apnea and its response to acute CPAP treatment. *PloS one.* 2012;7:e33769. <https://doi.org/10.1371/journal.pone.0033769>