

## TRANSIENT UPREGULATION OF PROTEIN KINASE C IN PRESSURE-OVERLOADED NEONATAL RAT MYOCARDIUM

**B. Hamplová<sup>1,2</sup>, F. Novák<sup>2,4</sup>, F. Kolář<sup>3,4</sup>, O. Nováková<sup>1</sup>**

*<sup>1</sup>Department of Cell Biology, <sup>2</sup>Department of Biochemistry, Charles University, Faculty of Science, <sup>3</sup>Institute of Physiology, Academy of Sciences of the Czech Republic and <sup>4</sup>Centre for Cardiovascular Research, Prague, Czech Republic.*

*Address for offprints:*

O. Nováková,

Charles University in Prague, Faculty of Science,

Department of Cell Biology, Viničná 7, 128 44, Prague 2, Czech Republic

Fax: +420 2 21951758

E-mail: [olnov@natur.cuni.cz](mailto:olnov@natur.cuni.cz)

Short title: Protein kinase C in pressure-overloaded neonatal myocardium

## Summary

Protein kinase C (PKC) appears to play a significant role in the signal transduction of cardiac growth and development. The aim of this study was to determine changes in the total PKC activity and the expression of PKC isoforms  $\alpha$ ,  $\delta$  and  $\epsilon$  in the rat heart that was affected by pressure overload imposed at postnatal day (d) 2. Three groups of Wistar rats were employed for the experiment: rats submitted to the abdominal aortic constriction (AC), sham-operated controls (SO) and intact controls. Animals were sacrificed at d2, 3, 5 and 10. The total PKC activity was measured by the incorporation of  $^{32}\text{P}$  into histone H1S and the expression of PKC was analyzed by immunoblotting in the homogenate of the left ventricular myocardium and in the cytosolic, membrane-enriched ( $10^5 \times \text{g}$ ) and nuclear-cytoskeletal-myofilament-enriched ( $10^3 \times \text{g}$ ) fractions. We observed the significant transient increase in both the total PKC activity and the expression of all isoforms at d5 (the 3<sup>rd</sup> day after the operation) in the cardiac homogenate of AC rats as compared with SO animals. Aortic constriction did not significantly affect the distribution of activity and isoform abundance among individual cellular fractions except for PKC $\delta$ , which increased significantly at d10 in the cytosolic fraction at the expense of the membrane-enriched fraction. It is concluded that PKC $\alpha$ , PKC $\delta$  and PKC $\epsilon$  undergo transient upregulation associated with the accelerated cardiac growth induced by pressure overload imposed in the very early postnatal period.

*Key words:* protein kinase C, pressure overload, postnatal development, rat myocardium

## Introduction

The developing mammalian heart has to overcome an increased workload, which is associated with the rapid growth of the left ventricle during postnatal ontogeny. Just after birth, the myocardium continues growing by adding new cells (hyperplasia), but the further growth of the heart is due to the increase of cell volume (hypertrophy) as ventricular cardiomyocytes lose their capability of dividing soon (Clubb and Bishop 1984, Li et al. 1996), though limited proliferative activity may restore under certain conditions even in adulthood (Beltrami et al. 2001, Bruel et al. 2007). It is well known that protein kinase C (PKC) is an important player in the complex regulation of cell proliferation and myocardial hypertrophy (Bogoyevitch and Sugden 1996, Sabri and Steinberg 2003, Buchner 2000). The Gq/PLC/PKC signaling pathway was proved to mediate both the normal cardiac postnatal growth and hypertrophic growth of pressure-overloaded hearts (Dorn and Force 2005, Wettschureck et al. 2001).

PKC consists of a family of at least 11 isozymes, which are divided into three groups. The members of a classical group ( $\alpha$ ,  $\beta_{1/2}$  and  $\gamma$ ) are  $\text{Ca}^{2+}$  dependent and need 1,2-diacylglycerol (DAG) and phosphatidylserine (PS) for their activation; novel PKC isoforms ( $\delta$ ,  $\epsilon$ ,  $\eta$  and  $\theta$ ) are  $\text{Ca}^{2+}$  independent and are activated by DAG and PS; atypical isoforms ( $\zeta$  and  $\iota/\lambda$ ) are  $\text{Ca}^{2+}$  and DAG independent but require PS as a cofactor. PKC isozymes are expressed differentially in various organs and tissues (Wetsel et al. 1992). PKC $\alpha$ , PKC $\delta$ , PKC $\epsilon$  and PKC $\zeta$  were mostly identified in a neonatal rat heart and all of them diminished markedly until adulthood (Clerk et al. 1995, Rybin and Steinberg 1994, Hamplova et al. 2005). However, we have recently described rapid day-by-day developmental changes in the expression of PKC $\alpha$ , PKC $\delta$  and PKC $\epsilon$  in the homogenate and the cellular fractions of rat heart during the first 10 postnatal days. We observed more than 2-fold transient increase in the content of PKC $\delta$  and PKC $\epsilon$  associated with nuclear-cytoskeletal-myofilament structures

between d3 and 5 (Hamplova et al. 2005). It is worth noting that this narrow developmental period is connected with the rapid transition from the hyperplastic to the hypertrophic growth (Li et al. 1996).

In the present study we were interested to find out whether the aortic constriction, an additional growth-promoting stimulus targeted just prior to period of hyperplastic-to-hypertrophic transition, can influence the developmental changes in the total activity and the expression and subcellular distribution of PKC $\alpha$ , PKC $\delta$  and PKC $\epsilon$ . We used the experimental model of the neonatal pressure-overloaded heart that allows observing a response of the myocardium that is still in the proliferative phase of the cardiomyocyte growth (Kolar et al. 1998, Sedmera et al. 2003).

## **Methods**

### *Animal model*

Newborn male Wistar rats were obtained from the animal care facility of the Institute of Physiology, Prague, Czech Republic. Under light ether anesthesia, pressure overload was induced by the constriction of the abdominal aorta (AC) at postnatal day 2 as described earlier (Kolar et al. 1998). Sham-operated (SO) littermates were used as the age-matched controls; their aorta was exposed, but not constricted. Intact littermates were also used. **The animals were sacrificed at d2 (intact subgroup only), d3, d5 and d10.** Their hearts were dissected free of an atrial tissue, large blood vessels and the right ventricle. The left ventricle including the septum (LV+S) was rinsed in cold saline (5 °C), weighed, frozen in liquid nitrogen and stored at –80 °C until use. This study conforms to the *Guide for the Care and Use of Laboratory Animals* published by the U.S. National Institute of Health (National Institutes of Health publication No. 85-23, revised 1996).

### *Tissue fractionation*

A specimen for a fractionation contained ventricles from 4 (d10) to 14 (d2) animals. The tissue was minced using Turrax (30 s, twice) and then homogenized in 10 volumes of ice-cold homogenization buffer composed of (in mmol/l): 12.5 Tris-HCl (pH 7.4), 250 sucrose, 2.5 EGTA, 1 EDTA, 100 NaF, 5 dithiothreitol, 0.3 phenylmethylsulfonyl fluoride, 0.2 leupeptin, 0.02 aprotinin. The homogenate was fractionated according to the method described previously (Gu and Bishop 1994), with slight modifications. Briefly, the homogenate was centrifuged at  $100 \times g$  for 20 min to remove cellular debris and unbroken cells. A supernatant was centrifuged at  $1000 \times g$  for 10 min to produce a nuclear-cytoskeletal-myofilament-enriched fraction (denoted as *nuclear fraction* in the text for simplicity) followed by centrifugation at  $100,000 \times g$  for 60 min. The  $100,000 \times g$  pellet contained a membrane-enriched fraction and the supernatant was a cytosolic fraction. The homogenate and the pellets of nuclear and membrane fractions were resuspended in the homogenization buffer containing 1 % Triton X-100, held on ice for 60 min and centrifuged at  $100,000 \times g$  for further 60 min. The resulting detergent-treated supernatants were used for activity and immunoblotting analyses. Triton X-100 was also added to the cytosolic fraction to give the final concentration of 1 %. Protein content was determined according to Lowry, as modified by Peterson (1977).

### *Activity assay of total PKC*

The PKC activity was determined as a difference in the incorporation of  $^{32}P$  from  $\gamma[^{32}P]ATP$  into a histone H1S in the presence and the absence of PS, 1-octanoyl-2-acetyl-glycerol (OAG) and  $Ca^{2+}$ . The required amount of PS and OAG dissolved in chloroform was dried in the stream of nitrogen and solubilized in 0.3 % Triton X-100 by vortexing and incubation at 30 °C for 5 min. A reaction mixture (0.1 ml) contained (in mmol/l): 20 Tris-HCl (pH 7.45), 10  $MgCl_2$ , 1 dithiothreitol, 1  $CaCl_2$ , 0.1 vanadate, 21  $\mu g$  PS, 5.1  $\mu g$  OAG and protein (in  $\mu g$ : 2.5 homogenate, 3 cytosolic, 2 nuclear, 1 membrane fractions). The reaction was started by the

addition of 10  $\mu$ l of 1 mM  $\gamma$ [<sup>32</sup>P]ATP (50-100 cpm/pmol) and terminated after 5 min at 30 °C by the addition of 1 ml ice-cold 25 % trichloroacetic acid. The resulting precipitates were filtered through pre-soaked nitrocellulose filters (Pragopor, Pragochema) and washed three times with 5 % trichloroacetic acid. The radioactivity was quantified using the Cerenkov decay.

#### *Immunoblot analyses of PKC isoforms*

Samples were electrophoresed on the 8 % bis-acrylamide polyacrylamide gel. SDS-PAGE was carried out at 20 mA/gel for 90 min on a Mini-Protean II apparatus (Bio-Rad). After the electrophoresis, resolved proteins were transferred to a nitrocellulose membrane (Amersham International). The membranes were incubated in 5 % dry low-fat milk in Tris-buffered saline with Tween 20 (TTBS) for 60 min at room temperature in order to block a nonspecific binding. After washing in TTBS buffer (3-times quickly, 3-times 5 min each) they were probed with the PKC isoform-specific primary rabbit antisera (1:8000 in TTBS) for 90 min at room temperature. The membranes were washed again and incubated with the secondary swine anti-rabbit IgG antibody labeled with the horseradish peroxidase (1:4000 in TTBS) for 60 min at room temperature. Before an enhanced chemiluminescence (ECL), the nitrocelluloses were washed as described above and stored in TTBS for at least 2 hours. For ECL, we prepared ECL substrates A (Luminol solution) and B (H<sub>2</sub>O<sub>2</sub> solution); they were mixed 1:1 and poured on the immunoblot. A specific signal was documented on an autoradiography film (Amersham Int). A scanning (Epson Perfection 1240U) and ImageQuant software were used for the quantification of a relative abundance of the individual PKC isoforms. To ensure the specificity of the immunoreactive proteins, Western blots were done in the presence and the absence of competing immunizing peptides (see Fig. 1 in Hamplova et al. 2005). The PKC  $\delta$  isoform was identified as the doublet of proteins. Both proteins were recognized using the antibody mentioned above, blocked by the appropriate immunizing

peptide and taken for the quantification in our study. Similarly, Ogita *et al.* (1992) immunodetected PKC  $\delta$  as the doublet of proteins (78 and 76 kDa) in the preparation from rat brain; the 78 kDa protein was identified as the phosphorylated form of the 76 kDa protein by means of the protein phosphatase 2A treatment.

### *Chemicals*

Chemical reagents including the PKC isoform-specific primary antisera and competing immunizing peptides were obtained from Sigma-Aldrich. The secondary swine anti-rabbit IgG antibody labeled with a horseradish peroxidase was from Sevapharma. Radioactive  $\gamma$ [<sup>32</sup>P]ATP was obtained from ICN Biomedicals.

### *Statistical analyses*

All results are expressed as means  $\pm$  S.E.M. The statistical significance of differences was determined by one-way ANOVA and subsequent Newman-Keuls test ( $p < 0.05$ ).

## **Results**

### *Weight parameters*

Body weight was significantly altered by neither SO, nor AC in any age group of the animals. AC led to the mild left ventricle enlargement, as evident from increases of LV+S/BW ratio by 14% and 23% at d5 and d10 (i.e. 3 and 8 days after the surgery), respectively, as compared with the SO littermates (Table 1).

### *Total PKC activity*

The total PKC activity in the homogenate of the intact ventricular myocardium during early postnatal development, and the effect of SO and AC on the total activity within the period of the study are shown in Fig. 1A. The PKC activity declined gradually from d2 till d10 in intact

rats. SO did not induce any changes. AC caused the transient increase in the activity (by 45%) at d5 as compared with the SO rats. The changes after SO and AC in the total activity in cellular fractions are presented in Fig. 1B. Compared to intact controls, we observed relative increase in the PKC activity at d3 in the nuclear fraction of SO group that occurred at the expense of the membrane and the cytosolic fractions.

#### *PKC isoforms expression and distribution*

The PKC $\alpha$  expression declined gradually from d2 till d10 in the homogenate of the intact myocardium. SO influenced the expression of PKC $\alpha$  neither in the homogenate nor in any cellular fraction. AC transiently increased the expression of PKC $\alpha$  by 55% at d5 in the homogenate but did not affect its distribution between fractions (Fig. 2).

The PKC $\delta$  expression declined from d2 till d10 in the myocardial homogenate of intact animals. SO had no effect while AC transiently elevated the expression of PKC $\delta$  by 67% at d5 as compared with the SO group. As for the distribution of PKC $\delta$ , we observed the relative decrease in the nuclear fraction at d5 by 13% and the increase in the membrane fraction at d5 by 8% and at d10 by 11% in SO rats as compared with intact controls. In the group of AC rats, we found the relative decrease of  $\delta$  isoform in the membrane fraction (by 13%) and the increase in cytosol at d10 in the comparison with SO animals (Fig. 3).

The expression of PKC $\epsilon$  declined from d2 till d10 with the sharpest drop by d3 in the cardiac homogenate of intact rats. SO had no effect on PKC $\epsilon$  expression and AC transiently increased it at d5 by 39% as compared with SO controls. Neither SO nor AC influenced the distribution of PKC $\epsilon$  among cellular fractions (Fig.4).



## **Discussion**

The aim of this study was to examine the effect of pressure overload on the pattern of developmental changes in the total PKC activity and the expression of PKC $\alpha$ , PKC $\delta$  and PKC $\epsilon$  in the rat heart during the first ten postnatal days. Although several studies investigated the role of the PKC family in the regulation of the cardiac hypertrophy induced by the aortic constriction (Gu and Bishop 1994, Bayer et al. 2003, Braun et al. 2002), none of them examined the cardiomegaly induced before d5, i.e. in the proliferative phase of the rat cardiomyocyte growth (Li et al. 1996). It should be stressed that the myocardium affected at this time of development responds differently to pressure overload as compared with the adult heart. When the aortic constriction was performed early after birth (d2), the increase of the left ventricular mass resulted first from the stimulated cellular hyperplasia followed later by the cardiomyocytes enlargement (Sedmera et al. 2003). Moreover, cardiomegaly was associated with capillary proliferation in proportion to the increase of ventricular mass in this experimental model, and exhibited neither increased rates of apoptosis nor myocardial fibrosis characteristic of adult pressure-overloaded hearts (Kolar et al. 1998, Sedmera et al. 2003, Oliviero et al. 2000). Thus, aortic constriction in neonates results in a unique cardiac phenotype that markedly differs from that induced in the adulthood, and accordingly the PKC isoforms might be differently involved in its regulation.

In adult pressure-overloaded myocardium, PKC $\alpha$  and PKC $\delta$  were markedly upregulated, the latter one even as early as the first day after aortic constriction (Bayer et al. 2003, Braun et al. 2002). These results provide indirect evidence that PKC $\delta$  might be involved in the induction of compensatory hypertrophy in the adult heart after aortic constriction, whereas both PKC $\alpha$  and PKC $\delta$  might be responsible for the transition to heart failure. The studies with transgenic mice show that the function of PKC $\delta$  depends on the strength of hypertrophic stimuli (Hahn et al. 2002). A modest increase in PKC $\delta$  expression was shown to

result in so called “physiological” hypertrophy, but on the other hand, high chronic expression of PKC $\delta$  evoked cardiomyocyte necrosis and contractile dysfunction (Chen et al. 2001, Hahn et al. 2002). The role of PKC $\epsilon$  in the regulation of cardiac hypertrophy appears to be parallel with that of PKC $\delta$  during compensatory hypertrophy (Chen et al. 2001, Mochly-Rosen et al. 2000, Takeishi et al. 2000). Besides, Heidkamp et al. (2001) showed that just PKC $\delta$  and PKC $\epsilon$  act as a switch between cardiomyocyte hypertrophy and apoptosis, and PKC $\epsilon$  selectively activates the MAPK cascade implicated in growth response and cell survival, whereas PKC $\delta$  preferentially activates stress-activated protein kinase cascades implicated in detrimental changes of the heart. Thus, it seems that the fate of cardiomyocytes in pressure-overloaded hearts depends mostly on the balance of nPKC isoforms to downstream signaling cascades.

In this study, we have found the transient increase in PKC activity and upregulation of all isoforms (PKC $\alpha$ , PKC $\delta$  and PKC $\epsilon$ ) at d5 (it means the 3<sup>rd</sup> day after the operation) in the myocardium of AC rats. Already at this early stage, the slight but significant increase in the relative left ventricular mass was observed. It can be speculated that the delay in the normal postnatal decline in PKC activity/expression caused by aortic constriction is related to **transitory stimulation of myocyte hyperplasia and a delayed transition to hypertrophic growth.** As for the redistribution of the PKC isoforms between cellular fractions, we did not observe any major changes except for the decreased proportion of PKC $\delta$  in the membrane fraction and, conversely, the increased proportion of this isoform in the cytosol only at d10 in hearts of pressure-overloaded rats. Noteworthy, PKC $\alpha$  and PKC $\epsilon$  had a tendency to copy these changes in PKC $\delta$  distribution. **Altogether, these data suggest that individual PKC isoforms may act in synergy during the early phase of pressure overload in this model. However, it is preliminary to speculate about possible implication of these changes for an accelerated myocardial growth response.**

PKC isoforms can be activated by any membrane receptor that couples to Gq. It is unknown which of these receptors contribute to PKC activation and acceleration of cardiac growth in our experimental model of neonatal pressure-overloaded hearts. The only data obtained on the same model concern angiotensin II (Ang II) and its receptor types AT1 and AT2. It was demonstrated that Ang II content in sera increased transiently already 2 h after aortic constriction. At the receptor level, a transient increase in both AT1 and AT2 was observed one day after the surgery. Moreover, AT2 transcript levels were linearly correlated with the left ventricular mass in the AC group (Oliviero et al. 2000). The rise in Ang II level preceding the increase in Ang II binding capacity and PKC activation/upregulation might suggest that this peptide is one of the triggers implicated in the PKC-mediated early response of neonatal myocardium to pressure overload. Interestingly, in the normal rat heart, the expression of AT receptors is maximal just after birth (Hunt et al. 1995) decreasing rapidly thereafter together with PKC activity and isoform expression (Hamplova et al. 2005). However, these correlative studies do not allow to make any clear conclusion because PKC is not the only signaling pathway activated by Ang II and a variety of other factors might also contribute to accelerated neonatal cardiac growth in response to pressure overload, either mediated by PKC or not.

While the operation itself did not influence the total PKC activity and isoforms expression in the cardiac homogenate, it is necessary to note that it had significant effects on distributions of PKC activity and PKC $\delta$  abundance among cellular fractions compared with intact littermates. This finding is in line with the view that PKC may play a role in myocardial inflammatory response to stress and injury. For example, Tan et al. (2007) demonstrated a marked increase in PKC $\alpha$ , PKC $\delta$  and PKC $\epsilon$  abundance in membranes and PKC-dependent secretion of pro-inflammatory cytokines by cardiomyocytes induced by burn injury. In our study, only PKC $\delta$  redistributed to the membrane fraction in the SO group suggesting that the

surgical intervention itself was relatively mild stimulus. Moreover, an effect of anesthesia on PKC distribution cannot be excluded.

In conclusion, we demonstrated a transient increase in PKC activity and isoform expression at d5 in the rat ventricular myocardium subjected to pressure overload induced by aortic constriction at postnatal day 2. These effects were not accompanied by major changes in subcellular distribution of individual PKC isoforms. Although it is likely that PKC signaling is involved in mediating the early hyperplastic/hypertrophic myocardial growth response, further studies are needed to prove this hypothesis.

### **Acknowledgements**

The authors thank J. Ševčíková and M. Pešková for their excellent technical assistance. This work was supported by grant MSM 0021620858.

## References

- BAYER AL, HEIDKAMP MC, PATEL N, PORTER M, ENGMAN S, SAMAREL AM: Alterations in protein kinase C isoenzyme expression and autophosphorylation during the progression of pressure overload-induced left ventricular hypertrophy. *Mol Cell Biochem* **242**: 145-152, 2003.
- BELTRAMI AP, URBANEK K, KAJSTURA J, YAN S-M, FINATO N, BUSSANI R, NADAL-GINARD B, SILVESTRI F, LERI A, BELTRAMI A, ANVERSA P: Evidence that human cardiac myocytes divide after myocardial infarction. *N Engl J Med* **344**: 1750-1757, 2001.
- BOGOYEVITCH MA, SUGDEN PH: The role of protein kinases in adaptational growth of the heart. *Int J Biochem Cell Biol* **28**: 1-12, 1996.
- BRAUN MU, LAROSEE P, SCHON S, BORST MM, STRASSER RH: Differential regulation of cardiac protein kinase C isozyme expression after aortic banding in rat. *Cardiovasc Res* **56**: 52-63, 2002.
- BRÚEL A, CHRISTOFFERSEN TEH, NYENGAARD JR: Growth hormone increases the proliferation of existing cardiac myocytes and the total number of cardiac myocytes in the rat heart. *Cardiovasc Res* **76**: 400-408, 2007.
- BUCHNER K: The role of protein kinase C in the regulation of cell growth and in signalling to the cell nucleus. *J Cancer Res Clin Oncol* **126**: 1-11, 2000.
- CHEN L, HAHN H, WU G, CHEN CH, LIRON T, SCHECHTMAN D, CAVALLARO G, BANCIL, GUO Y, BOLLI R, DORN GW, MOCHLY-ROSEN D: Opposing cardioprotective actions and parallel hypertrophic effects of delta PKC and epsilon PKC. *Proc Natl Acad Sci USA* **98**: 11114-11119, 2001.

CLERK A, BOGOYEVITCH MA, FULLER SJ, LAZOU A, PARKER PJ, SUGDEN PH. Expression of protein kinase C isoforms during cardiac ventricular development. *Am J Physiol* **269**: H1087-H1097, 1995.

CLUBB FJJ, BISHOP SP: Formation of binucleated myocardial cells in the neonatal rat. An index for growth hypertrophy. *Lab Invest* **50**: 571-577, 1984.

DORN GW, FORCE T: Protein kinase cascades in the regulation of cardiac hypertrophy. *J Clin Invest* **115**: 527-537, 2005.

GU X, BISHOP SP: Increased protein kinase C and isozyme redistribution in pressure-overload cardiac hypertrophy in the rat. *Circ Res* **75**: 926-931, 1994.

HAHN HS, YUSSMAN MG, TOYOKAWA T, MARREEZ Y, BARRETT TJ, HILTY KC, OSINSKA H, ROBBINS J, DORN GW: Ischemic protection and myofibrillar cardiomyopathy: dose-dependent effects of in vivo deltaPKC inhibition. *Circ Res* **91**: 741-748, 2002.

HAMPLOVA B, NOVAKOVA O, TVRZICKA E, KOLAR F, NOVAK F: Protein kinase C activity and isoform expression during early postnatal development of rat myocardium. *Cell Biochem Biophys* **43**: 105-117, 2005.

HEIDKAMP MC, BAYER AL, MARTIN JL, SAMAREL AM: Differential activation of mitogen-activated protein kinase cascades and apoptosis by protein kinase C epsilon and delta in neonatal rat ventricular myocytes. *Circ Res* **89**: 882-890, 2001.

HUNT RA, CIUFFO GM, SAAVEDRA JM, TUCKER DC: Quantification and localisation of angiotensin II receptors and angiotensin converting enzyme in the developing rat heart. *Cardiovasc Res* **29**: 834-840, 1995.

KOLAR F, PAPOUSEK F, PELOUCH V, OSTADAL B, RAKUSAN K: Pressure overload induced in newborn rats: effects on left ventricular growth, morphology, and function. *Pediatric Res* **43**: 521-526, 1998.

LI F, WANG X, CAPASSO JM, GERDES AM: Rapid transition of cardiac myocytes from hyperplasia to hypertrophy during postnatal development. *J Mol Cell Cardiol* **28**: 1737-1746, 1996.

MOCHLY-ROSEN D, WU G, HAHN H, OSINSKA H, LIRON T, LORENZ JN, YATANI A, ROBBINS J, DORN GW: Cardiostrophic effects of protein kinase C epsilon: analysis by in vivo modulation of PKCepsilon translocation. *Circ Res* **86**: 1173-1179, 2000.

OGITA K, MIYAMOTO S, YAMAGUCHI K, KOIDE H, FUJISAWA N, KIKKAWA U, SAHARA S, FUKAMI Y, NISHIZUKA Y: Isolation and characterization of delta-subspecies of protein kinase C from rat brain. *Proc Natl Acad Sci USA* **89**: 1592-1596, 1992.

OLIVIERO PF, CHASSAGNE CF, KOLAR F, ADAMY CF, MAROTTE FF, SAMUEL JL, RAPPAPORT LF, OSTADAL B: Effect of pressure overload on angiotensin receptor expression in the rat heart during early postnatal life. *J Mol Cell Cardiol* **32**: 1631-1645, 2000.

PETERSON GL: A simplification of the protein assay method of Lowry et al. which is more generally applicable. *Anal Biochem* **83**: 346-356, 1977.

RYBIN VO, STEINBERG SF: Protein kinase C isoform expression and regulation in the developing rat heart. *Circ Res* **74**: 299-309, 1994.

SABRI A, STEINBERG SF: Protein kinase C isoform-selective signals that lead to cardiac hypertrophy and the progression of heart failure. *Mol Cell Biochem* **251**: 97-101, 2003.

SEDMERA D, THOMPSON RP, KOLAR F: Effect of increased pressure loading on heart growth in neonatal rats. *J Mol Cell Cardiol* **35**: 301-309, 2003.

TAKEISHI Y, PING P, BOLLI R, KIRKPATRICK DL, HOIT BD, WALSH RA: Transgenic overexpression of constitutively active protein kinase C epsilon causes concentric cardiac hypertrophy. *Circ Res* **86**: 1218-1223, 2000.

TAN J, MAASS DL, WHITE J, HORTON JW: Effects of burn injury on myocardial signaling and cytokine secretion: possible role of PKC. *Am J Physiol Regul Integr Comp Physiol* **292**: R887-R896, 2007.

WETSEL WC, KHAN WA, MERCHENTHALER I, RIVERA H, HALPERN AE, PHUNG HM, NEGRO-VILAR A, HANNUN YA: Tissue and cellular distribution of the extended family of protein kinase C isoenzymes. *J Cell Biol* **117**: 121-133, 1992.

WETTSCHURECK N, RUTTEN H, ZYWIETZ A, GEHRING D, WILKIE TM, CHEN J, CHIEN KR, OFFERMANN S: Absence of pressure overload induced myocardial hypertrophy after conditional inactivation of Galphaq/Galpha11 in cardiomyocytes. *Nat Med* **7**: 1236-1240, 2001.



Table 1. Weight parameters of intact, sham-operated and aortic-constricted rats during early postnatal development.

Postnatal days	n	BW (g)	LV+S (mg)	RV (mg)	LV+S/BW (mg/g)
2 day intact	44	6.61 ± 0.40	21.89 ± 0.46	7.83 ± 0.26	3.22 ± 0.05
3 day intact	24	9.04 ± 0.49	30.11 ± 0.88	9.64 ± 0.30	3.43 ± 0.10
SO	21	9.44 ± 0.98	32.92 ± 1.15	9.78 ± 0.55	3.40 ± 0.05
AC	21	7.45 ± 0.50	32.44 ± 1.63	10.14 ± 0.51	3.56 ± 0.07
5 day intact	24	10.78 ± 0.97	31.56 ± 1.29	10.10 ± 0.53	3.03 ± 0.05
SO	18	14.52 ± 2.02	37.10 ± 1.86 *	13.04 ± 0.76 *	3.02 ± 0.04
AC	18	14.01 ± 1.00	42.32 ± 1.77	12.45 ± 0.61	3.45 ± 0.07 #
10 day intact	12	22.42 ± 0.98	61.23 ± 2.59	21.51 ± 1.09	2.71 ± 0.06
SO	12	23.71 ± 1.63	61.56 ± 2.50	21.13 ± 0.93	2.78 ± 0.05
AC	12	23.67 ± 2.03	72.02 ± 3.01 #	20.56 ± 1.04	3.42 ± 0.12 #

Values are means ± S.E.M. BW, body weight; LV, left ventricle; S, septum; RV, right ventricle; SO, sham operation; AC, aortic constriction; n, number of animals. \*P<0.05, significant difference vs. intact controls; #P<0.05, significant difference vs. sham-operated controls.

## Figure legend

Fig. 1

The effect of sham operation and aortic constriction on developmental changes in the total protein kinase C activity (A) in the homogenate of left ventricular myocardium and (B) its distribution between cytosolic (white columns), membrane (dotted columns) and nuclear-cytoskeletal-myofilament fractions (crosshatched columns) during the first 10 postnatal days. Values are expressed (A) as arbitrary units (the sum of specific radioactivity measured on postnatal days 2, 3, 5 and 10 is equal to 1) or % of corresponding control values, and (B) as mol % (the sum of all cellular fraction equals 100%). Data are means  $\pm$  S.E.M. from 3 fractionations in each age group. <sup>+</sup>P<0.05, significant difference vs. day 2; \*P<0.05, significant difference vs. intact controls; <sup>#</sup>P<0.05, significant difference vs. sham-operated controls.

Fig. 2

The effect of sham operation and aortic constriction on developmental changes in the expression of protein kinase C $\alpha$  (A) in the homogenate of left ventricular myocardium (representative immuno-blots are shown) and (B) its distribution between cytosolic (white columns), membrane (dotted columns) and nuclear-cytoskeletal-myofilament fractions (crosshatched columns) during the first 10 postnatal days. Values are expressed (A) as arbitrary units (the sum of densitometry volumes measured on postnatal days 2, 3, 5 and 10 is equal to 1) or % of corresponding control values and (B) as mol % (the sum of all cellular fraction equals 100%). Data are means  $\pm$  S.E.M. from 3 fractionations in each age group. <sup>#</sup>P<0.05, significant difference vs. sham-operated controls.

Fig. 3


The effect of sham operation and aortic constriction on developmental changes in the expression of protein kinase C $\delta$  (A) in the homogenate of left ventricular myocardium (representative imuno-blot are shown) and (B) its distribution between cytosolic (white columns), membrane (dotted columns) and nuclear-cytoskeletal-myofilament fractions (crosshatched columns) during the first 10 postnatal days. Values are expressed (A) as arbitrary units (the sum of densitometry volumes measured on postnatal days 2, 3, 5 and 10 is equal to 1) or % of corresponding control values and (B) as mol % (the sum of all cellular fraction equals 100%). Data are means  $\pm$  S.E.M. from 3 fractionations in each age group. <sup>+</sup>P<0.05, significant difference vs. day 2; \*P<0.05, significant difference vs. intact controls; <sup>#</sup>P<0.05, significant difference vs. sham-operated controls.

Fig. 4

The effect of sham operation and aortic constriction on developmental changes in the expression of protein kinase C $\epsilon$  (A) in the homogenate of left ventricular myocardium (representative imuno-blot are shown) and (B) its distribution between cytosolic (white columns), membrane (dotted columns) and nuclear-cytoskeletal-myofilament fractions (crosshatched columns) during the first 10 postnatal days. Values are expressed (A) as arbitrary units (the sum of densitometry volumes measured on postnatal days 2, 3, 5 and 10 is equal to 1) or % of corresponding control values and (B) as mol % (the sum of all cellular fraction equals 100%). Data are means  $\pm$  S.E.M. from 3 fractionations in each age group. \*P<0.05, significant difference vs. day 2; <sup>#</sup>P<0.05, significant difference vs. sham-operated controls.

Fig. 1

A.



B.




Fig. 2





Fig. 3




Fig. 4

